


The listing of claims will replace all prior versions, and listings, of claims in the application:


Listing of Claims:

1. (Previously Presented) A process for preparing tetrahydropterin of the following formula



or a tetrahydropterin compound of said tetrahydropterin that is substituted at the 6-, or 7- or 6- and 7- position or positions,

comprising hydrogenating pterin of the following formula



or a pterin compound of said pterin that is monosubstituted at the 6-, or 7- or 6- and 7- position or positions,

with hydrogen in a polar reaction medium in the presence of a hydrogenation catalyst that is a metal complex that is soluble in the reaction medium, wherein the catalyst contains a ligand which is (i) triarylphosphine, (ii) tetramethylene phenylphosphine (iii) pentamethylene phenylphosphine, or (iv) a bidentate ligand with a tertiary amine group and a phosphine group or with two tertiary phosphine groups as complexing groups, wherein the bidentate ligands form together with a metal atom a five- to ten membered ring.


2. (Previously Presented) A process according to claim 1, wherein the polar reaction medium is an aqueous or alcoholic reaction medium.

3. (Previously Presented) A process according to claim 1, wherein the pterin compound is folic acid, a folic acid salt, a folic acid ester, a folic acid ester salt or a dihydro form thereof, with the proviso that in the event of using folic acid, a carboxylic acid thereof or a dihydro form thereof, the reaction medium is aqueous, and in the event of using a folic acid ester, a folic acid ester salt or a dihydro form thereof, the reaction medium is an alcohol.

4. (Previously Presented) A process according to claim 1, wherein the metal complex contains a chiral ligand.

5. (Previously Presented) A process according to claim 3, wherein the metal complex contains a chiral ligand.

6. (Previously Presented) A process according to claim 5, wherein the folic acid ester salt is of formula III and is in the form of a single enantiomer or a mixture of enantiomers of formula III,



in which

one of R_1 or R_2 is H, and the other one of R_1 or R_2 is a monovalent hydrocarbon radical or a hydrocarbon radical attached via a carbon atom in which one or more carbon atoms are each independently replaced by oxygen, sulfur, NH, -N=, or -N(C₁-C₄ Alkyl)-, or both R_1 and R_2 independently of one another represent a monovalent hydrocarbon radical or a hydrocarbon radical attached via a carbon atom in which one or more carbon atoms are each independently replaced by oxygen, sulfur, NH, -N=, or -N(C₁-C₄ Alkyl)-, HA stands for a monobasic to tribasic inorganic or organic acid, and x denotes an integer from 1 to 6 or a fractional number between 0 and 6.

7. (Previously Presented) A process according to claim 6, wherein HA is unsubstituted or substituted phenylsulphonic acid.

8. (Previously Presented) A process according to claim 1, wherein said process is carried out at a hydrogen pressure of 1 to 500 bars.

9. (Previously Presented) A process according to claim 1, wherein said process is carried out at a temperature is 0 to 150⁰ C.

10. (Previously Presented) A process according to claim 1, wherein the molar ratio of pterin or pterin compound to catalyst is 10 to 100,000.


11. (Previously Presented) A process according to claim 1, wherein the reaction medium is water or water in admixture with an organic solvent.

12. (Previously Presented) A process according to claim 2, wherein the alcoholic reaction medium is an alcohol, or an alcohol in admixture with an organic solvent.


13. (Previously Presented) A process according to claim 1, wherein the metal complex contains a d-8 metal.

14-28. (Cancelled)

29. (Previously Presented) A process for preparing tetrahydropterin of the following formula



or a tetrahydropterin compound of said tetrahydropterin that is substituted at the 6-, or 7- or 6- and 7- position or positions,
comprising hydrogenating pterin of the following formula



or a pterin compound of said pterin that is monosubstituted at the 6-, or 7- or 6- and 7- position or positions,

with hydrogen in alcohol or in alcohol in admixture with an organic solvent in the presence of a hydrogenation catalyst that is a metal complex that is soluble in the reaction medium.

30-32. (Cancelled)


33. (Previously Presented) A process according to claim 3, wherein the hydrogenation is carried out at elevated pressure.

34. (Previously Presented) A process according to claim 1, wherein the metal complex contains iridium, rhodium or ruthenium.

35-36. (Cancelled)


37-39. (Cancelled)

40. (Previously Presented) A process for preparing tetrahydropterin of the following formula



or a tetrahydropterin compound of said tetrahydropterin that is substituted at the 6-, or 7- or 6- and 7- position or positions,

comprising hydrogenating pterin of the following formula




or a pterin compound of said pterin that is monosubstituted at the 6-, or 7- or 6- and 7- position or positions,

with hydrogen in alcohol or in alcohol in admixture with an organic solvent in the presence of a hydrogenation catalyst that is a metal complex that is soluble in the reaction medium, wherein the pterin compound is folic acid, a folic acid salt, a folic acid ester, a folic acid ester salt or a dihydro form thereof, with the proviso that in the event of using folic acid, a carboxylic acid thereof or a dihydro form thereof, the reaction medium is aqueous, and in the event of using a folic acid ester, a folic acid ester salt or a dihydro form thereof, the reaction medium is an alcohol.

41-44. (Cancelled)

45. (Previously Presented) A process according to claim 1, wherein the pterin compound is a pterin that is substituted in the 6- position.

46. (Previously Presented) A process according to claim 1, wherein the pterin compound is of formula (A)



in which

R₁₀₁ is H or independently has the meaning of R₁₀₀, and

R₁₀₀ is an organic radical attached via a C, O or N atom and having 1 to 50 carbon atoms.


47. (Previously Presented) A process according to claim 46, wherein R₁₀₀ contains 1 to 30 carbon atoms and is not interrupted or is interrupted by one or more of -O-, -NH-, -N(C₁-C₄-alkyl)-, -C(O)-, -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)NH-, -NHC(O)-, -NHC(O)O-, -OC(O)NH-, -NHC(O)NH-, -C(O)N(C₁-C₄-alkyl)-, -N(C₁-C₄-alkyl)C(O)-, -N(C₁-C₄-alkyl)C(O)O-, -OC(O)N(C₁-C₄-alkyl)-, -N(C₁-C₄-alkyl)C(O)N(C₁-C₄-alkyl)-, and which is unsubstituted or is substituted with F, Cl, Br, -CN, -OCN, -NCO, -OH, -NH₂, -NHC₁-C₄-alkyl, -N(C₁-C₄-alkyl)₂, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-hydroxyalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -C(O)OH, -C(O)OM₁₀₀, -C(O)OC₁-C₄-alkyl, -C(O)NH₂, -C(O)NHC₁-C₄-alkyl, -C(O)N(C₁-C₄-alkyl)₂, R₁₀₂-C(O)O-, R₁₀₂-OC(O)O-, R₁₀₂-C(O)NH-, R₁₀₂-C(O)N(C₁-C₄-alkyl)-, R₁₀₂-NHC(O)NH-, R₁₀₃C(O)- or -CH(O), wherein

M₁₀₀ is Li, K, Na, NH₄⁺, or ammonium with 1 to 16 carbon atoms,

R₁₀₂ is C₁-C₈-alkyl, C₅- or C₆-cycloalkyl, phenyl or benzyl, and


R₁₀₃ is C₁-C₄-alkyl, phenyl or benzyl.

48. (Previously Presented) A process for preparing tetrahydropterin of the following formula




or a tetrahydropterin compound of said tetrahydropterin that is substituted at the 6-, or 7- or 6- and 7- position or positions,

comprising hydrogenating pterin of the following formula



or a pterin compound of said pterin that is monosubstituted at the 6-, or 7- or 6- and 7- position or positions,

with hydrogen in a polar reaction medium in the presence of a hydrogenation catalyst that is a metal complex that is soluble in the reaction medium of formula XLIV, XLIVa or XLIVb,



in which

Y stands for monoolefin ligands or a diene ligand;

X₇ represents an achiral or chiral ditertiary diphosphine, that forms a 5 to 7 membered ring with the metal atom Me₂ or Ru;

X₇ represents an achiral or chiral ligand that forms a 5 to 7 membered ring with the metal atom Me₂ or Ru, wherein said ligand contains two tertiary phosphine groups;

Me₂ denotes Ir(I) or Rh(I);

Z represents -Cl, -Br, or -I; and

A₂ is ClO₄⁻, CF₃SO₃⁻, CH₃SO₃⁻, HSO₄⁻, BF₄⁻, B(Phenyl)₄⁻, PF₆⁻, SbCl₆⁻, AsF₆⁻ or SbF₆⁻;

X₈ and X₉ are the same or different and have the meaning of Z or A₂, or X₈ has the meaning of Z or A₂ and X₉ stands for hydride.

49. (Previously Presented) A process according to claim 6, wherein R₁ and/or R₂ are, each independently,

pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, piperazinyl, pyrrolidinyl methyl, pyrrolidinyl ethyl, piperidinyl methyl, piperidinyl ethyl, morpholinyl methyl, morpholinyl ethyl, tetrahydropyranyl methyl, tetrahydropyranyl ethyl, piperazinyl methyl or piperazinyl ethyl.

50-57. (Cancelled)